

Announcement: LISP Project and Extra Credit Project is due Thursday December 1, 2011.

Reading Assignment: Read Chapters 10-12 in Nilsson

Homework

Homework Number 10: Problems 10.5, 10.6, 12.1, 12.2 in Nilsson due Tuesday November 1 in class.

Homework Number 11: Problems 13.2, 14.1, 14.3, 14.4 and 14.5 in Nilsson due Tuesday November 8 in class.

Homework Number 12: Problems 16.1, 16.2, 16.3, 16.4 16.11 in Nilsson due Tu. Nov. 15, in class

Homework Number 13: Due Tuesday November 22, in class

Computation Deduction.

Using **Resolution Refutation** deduce the following computation to obtain a value for the goal by drawing the Consistent Solution Graph for the goal and *prove its consistency*. Make sure your graph is clearly marked and it follows a complete strategy.

Facts:

F1: $\text{member}(X, \text{cons}(X, Y))$.

F2: $\text{subset}(\text{nil}, Z)$.

Rules:

R1: $\text{member}(X2, Y2) \rightarrow \text{member}(X2, \text{cons}(U, Y2))$.

R2: $\text{member}(X3, Y3) \wedge \text{subset}(Z3, Y3) \rightarrow \text{subset}(\text{cons}(X3, Z3), Y3)$.

Goal: $\text{subset}(\text{cons}(3, \text{cons}(2, \text{nil})), \text{cons}(1, \text{cons}(2, \text{cons}(3, \text{cons}(4, \text{nil}))))))$.

{Note: If you prefer, you may use the notation $\text{subset}([3,2],[1,2,3,4])$ or $\text{subset}((3\ 2),(1\ 2\ 3\ 4))$.}

Draw the graph, show the substitutions are consistent, and obtain the value of the goal.

OUTLINE Class #26

Outline

- Adversarial Searches
- Minimax Search
- Tic-Tac-Toe Example